


AD-A177 100 ON BILINEAR FORMS IN GAUSSIAN RANDOM VARIABLES TOEPLITZ 1/1  
MATRICES AND PARS. (U) NORTH CAROLINA UNIV AT CHAPEL  
HILL CENTER FOR STOCHASTIC PROC. F AVRAM NOV 86 TR-14  
UNCLASSIFIED AFOSR-TR-87-0099 F49628-85-C-0144 F/G 12/1 NL





MICROCOPY RESOLUTION TEST CHART  
NBS 1010-A, UNITED STATES GOVERNMENT PRINTING OFFICE

UNCLASSIFIED

AD-A177 100

②


SECURITY CLASSIFICATION

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) technical report No. 164		5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR. 87-0099</b>	
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6c. ADDRESS (City, State and ZIP Code) Center for Stochastic Processes, Statistics Department, Phillips Hall 039-A, Chapel Hill, NC 27514		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144	
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304
11. TITLE (Include Security Classification) On bilinear forms in Gaussian random variables, Toeplitz matrices and Parseval's relation		TASK NO. H5	WORK UNIT NO.
12. PERSONAL AUTHOR(S) Avram, F.			
13a. TYPE OF REPORT preprint	13b. TIME COVERED FROM 10/86 TO 9/87	14. DATE OF REPORT (Yr., Mo., Day) November 1986	15. PAGE COUNT 10
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR	
XXXXXXXXXXXXXXXXXX		Keywords: Toeplitz matrices; trace; singular values; cumulants, large deviations.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p><b>Abstract.</b> We improve a result of Szegő on the asymptotic behaviour of the trace of products of Toeplitz matrices.</p> <p>As an application, we improve also his result on the limiting behaviour of the bilinear forms</p> $B_n = \sum_{i,j=1}^n a_{i-j} X_i X_j,$ <p>where <math>X_i</math> is a stationary Gaussian sequence. A large deviations result is derived as well.</p> <p style="text-align: right;">X</p>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL MAJOR WOOD		22b. TELEPHONE NUMBER (Include Area Code) 703 767-5025	22c. OFFICE SYMBOL AFOSR/NM

## CENTER FOR STOCHASTIC PROCESSES

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina



On bilinear forms in Gaussian random variables,

Toeplitz matrices and Parseval's relation

by

Florin Avram

Technical Report No. 164

November 1986

Approved for public release;  
distribution unlimited.

Approved for release by NSA on 05-10-2014 pursuant to E.O. 13526  
UNCLASSIFIED//FORN DISSEM  
This document is the property of the NSA and is to be controlled, stored, handled, transmitted, and disposed of in accordance with NSA policy. It is not to be released, distributed, or otherwise made available to the public or other personnel without proper authorization. If this document is found in an unsecured location, it should be reported to the nearest NSA official. If this document is found in a secure location, it should be reported to the nearest NSA official. If this document is found in a secure location, it should be reported to the nearest NSA official.

# On bilinear forms in Gaussian random variables, Toeplitz matrices and Parseval's relation

FLORIN AVRAM

University of North Carolina  
Center for Stochastic Processes  
and  
Purdue University  
Mathematics Department

**Abstract.** We improve a result of Szegő on the asymptotic behaviour of the trace of products of Toeplitz matrices.

As an application, we improve also his result on the limiting behaviour of the bilinear forms

$$B_n = \sum_{i,j=1}^n a_{i-j} X_i X_j,$$

where  $X_i$  is a stationary Gaussian sequence. A large deviations result is derived as well.

## 1. Statement of Results

A. We study below the asymptotic behaviour of bilinear forms

$$(1.1) \quad B_n = \sum_{i,j=1}^n a_{i-j} X_i X_j$$

where  $X_i$  is a mean zero stationary Gaussian sequence.

This problem was first studied in the book of Grenander and Szegő, "Toeplitz matrices and their applications" (1958), as an application of their theory of the asymptotic behaviour of the trace of products of Toeplitz matrices.

Recently, there has been a renewed interest in this problem. See Fox and Taqqu (1983) and (1986) and Taniguchi (1986).

In Theorem 1 below we improve the results of Grenander and Szegő on the asymptotics of the trace of products of Toeplitz matrices. This theorem can be viewed also as a generalization of Parseval's relation. As a corollary of Theorem 1, we get a result which

---

**Keywords and Phrases:** Toeplitz matrices, trace, singular values, cumulants, large deviations.

**A.M.S. 1980 Subject Classifications:** Primary, 60F05; Secondary, 60F10.

This research supported by the Air Force Office of Scientific Research Contract No. F49620 85C 0144.

improves Theorem 11.6 of Grenander and Szegő on the bilinear forms  $B_n$  (See Theorem 2).

The proof of Theorem 1 is based on a norm inequality (See Theorem 3), communicated to us by Professor Larry Brown.

In a different direction, we establish a large deviations result about  $B_n$  (See Theorem 4).

B. Let:

$$(1.2) \quad r_n = EX_0 X_n$$

denote the covariance of the sequence  $X_n$ . The key fact about the bilinear form  $B_n$  is that its cumulants are:

$$(1.3) \quad \text{cum}_k(B_n) = 2^{k-1}(k-1)! \text{Tr}(A_n R_n)^k,$$

where  $A_n, R_n$  are the  $n \times n$  Toeplitz matrices:

$$A_n(i, j) = a_{i-j}, \quad R_n(i, j) = r_{i-j}, \quad \text{for } i, j = 1, \dots, n$$

(Formula 1.3) is an easy application of the "diagram" formula; see Rosenblatt (1985, Theorem 2.2)).

The first step in studying  $B_n$  should be thus the investigation of the asymptotic behaviour of the trace of products of Toeplitz matrices.

Let  $F_n^{(\nu)}$ ,  $\nu = 1, \dots, s$  be  $n \times n$  Toeplitz matrices of the form

$$F_n^{(\nu)}(i, j) = f_{i-j}^{(\nu)} \quad \text{for } i, j = 1, \dots, n \quad \text{and } \nu = 1, \dots, s,$$

and suppose  $f_k^{(\nu)}$  are the Fourier coefficients of the real, even functions  $f^{(\nu)}(x)$ , i.e.:

$$(1.4) \quad f_k^{(\nu)} = \int_{-\pi}^{\pi} e^{ikx} f^{(\nu)}(x) dx,$$

THEOREM 1. Suppose that

$$f^{(\nu)}(x) \in L_{p_\nu}, \quad 1 \leq p_\nu \leq \infty;$$

a) if  $\sum_{\nu=1}^s (p_\nu)^{-1} \leq 1$ , then

$$(1.5) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \text{Tr} \left( \prod_{\nu=1}^s F_n^{(\nu)} \right) = \int_{-\pi}^{\pi} \prod_{\nu=1}^s (2\pi f^{(\nu)}(x)) dx$$

b) if  $\alpha > 1$ , and  $\alpha \geq \sum_{\nu=1}^s (p_\nu)^{-1}$ , then

$$(1.6) \quad \lim_{n \rightarrow \infty} \frac{1}{n^\alpha} \text{Tr} \left( \prod_{\nu=1}^s F_n^{(\nu)} \right) = 0$$

Remarks: 1) Formula 1.5 was first obtained by Grenander and Szezö (1958), 7.4, under the assumption that  $f^{(\nu)}(x)$  are bounded.

2) Theorem 1a is also a generalization of the classical Parseval relation. Indeed, it is shown in the Appendix that the L.H.S. of 1.5 can also be written as the Caesaro sums:

$$(1.7) \quad \frac{1}{n} \text{Tr} \left( \prod_{\nu=1}^s F_n^{(\nu)} \right) = \frac{A_0 + \dots + A_{n-1}}{n},$$

where  $A_k$  are the "skew" convolution sums:

$$(1.8) \quad A_k = \sum_{\substack{\nu_1 + \dots + \nu_s = 0 \\ (\nu_1, \dots, \nu_s) \in D_k}} f_{\nu_1} \dots f_{\nu_s}^{(s)},$$

and

$$D_k = \{(\nu_1, \dots, \nu_s) : \max_{1 \leq j \leq s} \sum_{i=1}^j \nu_i - \min_{1 \leq j \leq s} \sum_{i=1}^j \nu_i \leq k\}.$$

Thus, Theorem 1a asserts the Cesaro convergence of the "skew" convolution sums.

Accepted for	
NTIS	
DTIC	
Unannounced	
Justification	
By	
Date	
A-1	
DTIC	
COPY	
INSPECTED	
6	

Note also that the "usual" convolution sums,

$$B_n = \sum_{\substack{j \in \{1, \dots, n\}^s \\ j_1 + \dots + j_s = 0}} f_{j_1}^{(1)} \dots f_{j_s}^{(s)}$$

converge to the R.H.S. of 1.5, if  $f^{(\nu)}(x) \in L_{p_\nu}$ ,  $\sum_{\nu=1}^s (p_\nu)^{-1} \leq 1$ , and  $1 < p_\nu < \infty$ , since then the Fourier sums of  $f^{(\nu)}(x)$  converge in  $L_{p_\nu}$  sense, and the scalar product is continuous. In this case, taking Cesaro sums is unnecessary. If, however, some  $p_\nu$  equal 1 or  $\infty$ , and  $\sum_{\nu=1}^s (p_\nu)^{-1} = 1$ , we do not know whether  $B_n$  converge to the R.H.S. of (1.5) in Cesaro sense. However, for  $n = 2$  and 3,  $C_n = D_n$ , and in the case  $n = 2$  we have the classical Parseval relation (See Katznelson, (1968), pg. 35).

As an immediate corollary of Theorem 1 we get:

**THEOREM 2.** Let  $a_k$  and  $r_k$  in 1.1) and 1.2) be the Fourier coefficients of the real even functions  $a(x)$  and  $r(x)$ , and suppose  $a(x) \in L_{p_1}$ ,  $r(x) \in L_{p_2}$ ,  $1 \leq p_1, p_2 \leq \infty$  and

$$(1.9) \quad (p_1)^{-1} + (p_2)^{-1} \leq 2^{-1}.$$

Then,

$$(1.10) \quad \frac{B_n - E(B_n)}{\sqrt{n}} \xrightarrow{d} N(0, \sigma^2),$$

where

$$\sigma^2 = 2(2\pi)^4 \int_{-\pi}^{\pi} a^2(x) r^2(x) dx.$$

*Proof:* Use the method of cumulants:

$$\text{cum}_k \left( \frac{B_n - EB_n}{\sqrt{n}} \right) = \begin{cases} 0 & \text{for } k = 1 \\ 2 \frac{\text{Tr}(A_n R_n)^2}{n} & \text{for } k = 2 \\ 2^{k-1} (k-1)! \frac{\text{Tr}(A_n R_n)^k}{n^{k/2}} & \text{for } k \geq 3 \end{cases}$$

$$\xrightarrow{n \rightarrow \infty} \begin{cases} 0 & \text{for } k = 1 \\ 2 \cdot (2\pi)^4 \int_{-\pi}^{\pi} a^2(x) r^2(x) dx & \text{for } k = 2, \text{ by Theorem 1a} \\ 0 & \text{for } k \geq 3, \text{ By Theorem 1b} \end{cases}$$



Notes: 1) 1.10 was first established by Grenander and Szegő (1958, Thm. 11.6), under the assumption that  $a(x)$  and  $r(x)$  are bounded.

2) Taqqu and Fox (1983) extended the result of Grenander and Szegő under a set of assumption different from ours. They show that if  $a(x)$  and  $r(x)$  are continuous, except maybe at 0, and are regularly varying at 0, then  $a(x)r(x) \in L_2$  (which is a weaker assumption than 1.9) is sufficient for 1.10 to hold.

C. Theorem 1 follows from the following inequality, communicated to us by Larry Brown:

THEOREM 3. For  $1 \leq p \leq \infty$ ,

$$(1.11) \quad \|F_n\|_p \leq n^{1/p} \|f(x)\|_p,$$

where  $\|F_n\|_p = (\sum_{j=1}^n |s_{j,n}|^p)^{1/p}$ ,  $s_{j,n}$  being the singular values of the matrix  $F_n$ .

(1.11) can be first established for  $p = 2, \infty$  and 1. By the Riesz convexity theorem, it follows then that it holds for every  $p$ .

D. We see from Theorem 1a that when  $a(x)$  and  $r(x)$  are bounded, the cumulants of  $B_n$  increase all at the same asymptotic rate ( $\text{cum}_k(B_n) = O(n)$ ). In such cases, large deviations results hold. We get, by applying Lemma 1 of Cox and Griffeath (1985), the following:

THEOREM 4. Suppose  $a(x)$  and  $r(x)$  are even, real functions, which are Riemann integrable. Let  $L = 4\pi \sup_x a(x) \cdot \sup_x r(x)$ , and  $\varphi(s) = -\frac{1}{2} \int_{-\pi}^{\pi} \ln(1 - 4\pi s a(x)r(x)) dx$ , for any  $s \in (-\infty, L^{-1})$ . Then,

a) for any  $\alpha \in (\varphi'(0), \lim_{s \nearrow L^{-1}} \varphi'(s))$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \Pr\left\{\frac{B_n}{n} > \alpha\right\} = -I(\alpha)$$

b) for any  $\alpha \in (\lim_{s \rightarrow -\infty} \varphi'(s), \varphi'(0))$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \Pr\left\{\frac{B_n}{n} < \alpha\right\} = -I(\alpha),$$

where  $I(\alpha) = \alpha s_\alpha - \varphi(s_\alpha)$ , and  $s_\alpha$  is the unique solution of  $\varphi'(s_\alpha) = \alpha$ .

## 2. Proofs

*Proof of Theorem 1:* a) Let  $m$  be the number of  $f^{(\nu)}$  which are non-polynomials (have infinitely many non zero Fourier coefficients). We will use induction on  $m$ . For  $m = 0$  (i.e. all  $f^{(\nu)}(x)$  are polynomials), it is easy to check that (1.5) holds. Suppose now (1.5) holds whenever we have at most  $m$  non-polynomials.

Consider then any set of  $f^{(\nu)}(x)$  which has at most  $m+1$  non-polynomials, and suppose w.l.o.g. that  $f^{(1)}(x)$  is a non-polynomial. Let then  $f_k^{(1)}(x)$  denote the  $k$ th Fejer sum of  $f^{(1)}(x)$ , let  $f^{(1),k}(x) = f^{(1)}(x) - f_k^{(1)}(x)$ , and let  $F_{n,k}^{(1)}$  and  $F_n^{(1),k}$  be the corresponding Toeplitz matrices. Then

$$(2.1) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \text{Tr}(F_{n,k}^{(1)} \prod_{\nu=2}^s F_n^{(\nu)}) = \int_{-\pi}^{\pi} 2\pi f_k^{(1)}(x) \prod_{\nu=2}^s (2\pi f^{(\nu)}(x)) dx$$

by the induction hypothesis, and the R.H.S. of (2.1) converges as  $k \rightarrow \infty$  to  $\int_{-\pi}^{\pi} \prod_{\nu=1}^s (2\pi f^{(\nu)}(x)) dx$  since  $1 \leq p_1 \leq \infty$  implies that  $\|f_k^{(1)} - f^{(1)}\|_{p_1} \xrightarrow{k \rightarrow \infty} 0$ , and  $\prod_{\nu=2}^s f^{(\nu)}(x) \in L_{q_1}$ , where  $(p_1)^{-1} + (q_1)^{-1} \leq 1$ . To show then that (1.5) holds with up to  $m+1$  non-polynomials it remains only to note that:

$$\begin{aligned} & \lim_{k \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} |\text{Tr} F_n^{(1),k} \prod_{\nu=2}^s F_n^{(\nu)}| \\ & \leq \lim_{k \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \|F_n^{(1),k} \prod_{\nu=2}^s F_n^{(\nu)}\|_1 \\ & \leq \lim_{k \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \frac{1}{n} \|F_n^{(1),k}\|_{p_1} \prod_{\nu=2}^s \|F_n^{(\nu)}\|_{p_\nu} \\ & \leq \lim_{k \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \frac{n \sum_{\nu} (p_\nu)^{-1}}{n} \|f^{(1),k}(x)\|_{p_1} \prod_{\nu=2}^s \|f^{(\nu)}(x)\|_{p_\nu} \quad (\text{by Thm 3}) \\ & = 0. \end{aligned} \quad \square$$

b) Assume first w.l.o.g.  $\sum_{\nu=1}^s (p_\nu)^{-1} > 1$ . (Otherwise the result follows from a)).

The proof is now similar with that of part a). If all  $f^{(\nu)}(x)$  are polynomials, the limit is 0 since  $\alpha > 1$ . Otherwise, if say,  $f^{(1)}(x)$  is a nonpolynomial, replace  $f^{(1)}$  by  $f^{(1),k} - f_k^{(1)}$ . Finally, let  $\theta = \sum_{\nu=1}^s (p_\nu)^{-1}$ , and note that

$$\begin{aligned} \frac{1}{n^\alpha} |Tr F_n^{(1),k} \prod_{\nu=2}^s F_n^{(\nu)}| &\leq \frac{1}{n^\alpha} \|F_n^{(1),k}\|_{\theta p_1} \prod_{\nu=2}^s \|F_n^{(\nu)}\|_{\theta p_\nu} \\ &\leq \frac{1}{n^\alpha} \|F_n^{(1),k}\|_{p_1} \prod_{\nu=2}^s \|F_n^{(\nu)}\|_{p_\nu} \quad (\text{since } \theta > 1) \\ &\leq \frac{1}{n^\alpha} \cdot n \sum_{\nu=1}^s (p_\nu)^{-1} \|f^{(1),k}\|_{p_1} \prod_{\nu=2}^s \|f^{(\nu)}\|_{p_\nu} \quad (\text{by Theorem 3}) \\ &\leq \|f^{(1),k}\|_{p_1} \prod_{\nu=2}^s \|f^{(\nu)}\|_{p_\nu} \xrightarrow{k \rightarrow \infty} 0. \quad \square \end{aligned}$$

*Proof of Theorem 4:* This is a straightforward application of Lemma 1 of Cox and Griffeath (1985). We need only to check that for any  $s \in (-\infty, L^{-1})$ , the cumulant generating function:

$\varphi_n(s) = \log E e^{s B_n}$  satisfies:

$$(2.2) \quad \lim_{n \rightarrow \infty} \varphi_n(s) = -\frac{1}{2} \int_{-\pi}^{\pi} \ell n(1 - 4\pi s a(x) r(x)) dx.$$

But  $\varphi_n(s)$  equals:

$$\varphi_n(s) = -\frac{1}{2} \sum_{i=1}^n \ell n(1 - 2s \lambda_{i,n}),$$

where  $\lambda_{i,n}$  are the eigenvalues of  $A_n, R_n$ , for any  $s \leq [\max_i 2\lambda_{i,n}]^{-1}$  (Direct computation).

(2.2) follows now Theorem 4.4 ii of Gray(1971), since  $a(x), r(x)$  are Riemann integrable, and the function  $\ell n(1 - 4\pi s z)$  is continuous for  $z \in (-\infty, \frac{L}{4\pi})$ , if  $s < L^{-1}$ .

□

**Note:** The assumption of Riemannian integrability is probably too strong. We follow however Gray in adopting it, due to the conceptual simplicity which it brings to the problem. (Under this assumption, the Toeplitz matrices are asymptotically equivalent

with circulant approximands, which are much easier to manipulate. This approach is nicely illustrated in Gray (1971)).

**Acknowledgement:** We thank Professor Larry Brown for communicating to us Theorem 3.

## Appendix

*Proof of formula 1.7:* Let

$$C_n = \{1, \dots, n\}^s,$$

let  $T(j_1, \dots, j_s)$  denote the range of sums  $\sum_{\nu=1}^k j_\nu$ , i.e.

$$T(j_1, \dots, j_s) = \text{Max}_{1 \leq k \leq s} \sum_{\nu=1}^k j_\nu - \text{Min}_{1 \leq k \leq s} \sum_{\nu=1}^k j_\nu,$$

let

$$D_n = \{(j_1, \dots, j_s) : \sum_{\nu=1}^s j_\nu = 0, T(j_1, \dots, j_s) \leq n\},$$

and let  $A_n$  be the "skew" convolution sums:

$$A_n = \sum_{\underline{j} \in D_n} f_{j_1}^{(1)} \dots f_{j_s}^{(s)}.$$

Then,

$$\begin{aligned} \frac{1}{n} T r \left( \prod_{\nu=1}^s F_n^{(\nu)} \right) &= \frac{1}{n} \sum_{\underline{i} \in C_n} f_{i_1 - i_2}^{(1)} f_{i_2 - i_3}^{(2)} \dots f_{i_s - i_1}^{(s)} \\ (A.1) \quad &= \frac{1}{n} \sum_{\underline{j} \in D_{n-1}} f_{j_1}^{(1)} \dots f_{j_s}^{(s)} \sum_{\substack{i_1 - i_2 = j_1 \\ i_s - i_1 = j_s}} 1 \\ &= \frac{1}{n} \sum_{\underline{j} \in D_{n-1}} f_{j_1}^{(1)} \dots f_{j_s}^{(s)} (n - T(j_1, \dots, j_s)). \end{aligned}$$

The last equality holds since the set of all  $\underline{i}$ 's with given  $\underline{j}$  differences can be obtained from any of its elements  $\underline{i}^{(0)}$ , by adding or subtracting  $(1, \dots, 1)$  as long as all components

are in the range  $\{1, \dots, n\}$ ; as such, it has  $(n - \max_{\nu} i_{\nu}^{(0)}) + \min_{\nu} i_{\nu}^{(0)}$  elements. Furthermore,

$$\begin{aligned} \max_{\nu} i_{\nu}^{(0)} - \min_{\nu} i_{\nu}^{(0)} &= \max_{\nu} (-i_{\nu}^{(0)}) - \min_{\nu} (-i_{\nu}^{(0)}) \\ &= \max_{\nu} (i_1^{(0)} - i_{\nu}^{(0)}) - \min_{\nu} (i_1^{(0)} - i_{\nu}^{(0)}) \\ &= \max_{\nu} \left( \sum_{k=1}^{\nu} j_k \right) - \min_{\nu} \left( \sum_{k=1}^{\nu} j_k \right) \\ &= T(j_1, \dots, j_s). \end{aligned}$$

Finally, from (A.1) we get

$$\frac{1}{n} \text{Tr} \left( \prod_{\nu=1}^s F_n^{\nu} \right) = \frac{1}{n} \sum_{k=0}^{n-1} \sum_{\underline{j} \in D_k} f_{j_1}^{(1)} \dots f_{j_s}^{(s)} = \frac{1}{n} \sum_{k=0}^{n-1} A_k.$$

## REFERENCES

- [1] FOX, R. AND TAQQU, M. S., *Central limit theorems for quadratic forms in random variables having long-range dependence*, School of Operations Research and Industrial Engineering Technical Report No.590 (1983), Cornell University.
- [2] —————, *Large-sample properties of parameter estimates for strongly dependent stationary Gaussian time series*, *Annals of Statistics* (1986), 517-532.
- [3] GRAY, R. M., *Toeplitz and circulant matrices*, Stanford Electron. Lab., Tech. Rep. 6502-1 (1971).
- [4] GRENANDER, V. AND SZEGÖ, G., *Toeplitz forms and their application*, Univ. California Press..
- [5] TANIGUCHI, M., *Berry-Esseen Theorems for Quadratic Forms of Gaussian Stationary Processes*, *Prob. Th. Rel. Fields* 1 **72** (1986), 185-194.
- [6] BROWN, LARRY, *private communication*.

END

4-~~2~~-87

DTIC